

Course No: DMED1209
Course Title: مهارات الإسعاف والطوارئ
Date: 26 /10/2017
No. of Questions: (3)
Time: One hour
Using Calculator (No)

University of Palestine

1st. Midterm First Aid
Exam
2017/2018
Total Grade:30 Marks

Instructor Name:
Student No.: _____
Student Name: _____
College Name: تخصص التمريض
Dep. / Specialist: _____
Using Dictionary (No)

QI. Read the following questions and choose the most correct answer (15 Marks):

1- All of the following aim of first aid:

- a) Preserve life.
- b) Prevent the situation from Worsening
- c) Promote the recovery.
- d) All of the above

2- When reaching an accident place what is the first thing a first aider should consider:

- a) Observing casualty skin color.
- b) Ensuring there is no danger.
- c) Seeing if the casualty is breathing.
- d) Obtaining help.

3- An unconscious casualty should be placed in the recovery position to:

- a) All the answer right
- b) Protect the airway and keep it open
- c) Keep them comfortable
- d) Prevent of developing shock

4- ABC stand for the following except:

- a) Airway, breathing, Cycle
- b) Apnea, breathing, Circulation
- c) Airway, back, Circulation
- d) Airway, breathing, Circulation

5- In primary survey for the victim primary survey include all of the following except :

- a) To make sure the Airway is open.
- b) To check they are breathing normally.
- c) Check for danger
- d) Check for major Bleeding and Burn.

6- All of the following are from the secondary survey for unconscious victim except

- a) Check for Bleeding
- b) Check for Burns
- c) Check for broken Bones.
- d) Check for Danger

7- When the casualty is not fully orientated but asks and answers your questions. We describe this condition of the casualty as

- a) fully oriented
- b) confused
- c) c -is fully alert
- d) The patient is able to speak words, but cannot put them together into logical sentences.

8- When you check for Danger, Response, Airway & Breathing this is called

- a) secondary survey
- b) tertiary survey
- c) primary survey
- d) all answers are correct

9- When you make compression to the adult causality, press down on the breastbone:

- a) 4 to 5cm
- b) 5 to 7cm
- c) 2 to 3cm
- d) 6 to 8cm.

10- Do 30 chest compressions at a rate of per minute:

- a) 70
- b) 100
- c) 80
- d) All are not true

11- Chest compressions for baby make by used:

- a) One hand
- b) Two hands
- c) Two fingers
- d) One fingers

12- Cardiopulmonary resuscitation cycle for adult, baby and child include:

- a) 30 compressions to 2 rescue breathing
- b) 15compressions to 2 rescue breathing
- c) 30 compressions to 1 rescue breathing
- d) 5 compressions to 1 rescue breathing

13- If there is more than one rescuer, change over every 2 minutes to

Course No: DMED1209

Course Title: مهارات الإسعاف والطوارئ

Date: 26 /10/2017

No. of Questions: (3)

Time: One hour

Using Calculator (No)

University of Palestine



1st. Midterm First Aid
Exam

2017/2018

Total Grade:30 Marks

Instructor Name:

Student No.: _____

Student Name: _____

College Name: تخصص التمريض

Dep. / Specialist: _____

Using Dictionary (No)

- a) Prevent infection
c) Prevent hemorrhage

- b) Prevent aspiration
d) Prevent fatigue

14- An interruption in the normal activity of the brain

- a) Consciousness
c) semiconscious

- b) unconsciousness
d) Disoriented

15- When check the breathing we do all of the following except

- a) Look.
c) Listen.

- b) Feel.
d) Compression.

Q2- Put (T) in front of the correct statement and (F) in front of the incorrect one. (15 Marks):

1	The first priority of first aid in emergency situation is to deal with major bleeding.	()
2-	Look, listen and feel for more than 20 seconds to check breathing.	()
3-	Compression and release shouldn't take an equal amount of time interval in CPR.	()
4-	If there is more than one rescuer in CPR, change over every 10 minutes to prevent fatigue.	()
5-	Unconsciousness can disable the body's natural reflexes such as coughing.	()
6	Secondary Survey should be done quickly & systematically.	()
7	When do Cardio Pulmonary Resuscitation (CPR) to the infant we do 2 rescue breathing first	()
8	When performing resuscitation on an infant, the rate of chest compressions to rescue breathing is 30 – 2	()
9	During (CPR) If the patient has vomited, turn them onto his side to prevent aspiration	()
10	The unconscious patient when laying on his side the tongue may fall back and blocking the airway	()

Course No: DMED1209
Course Title: مهارات الإسعاف والطوارئ
Date: 26 /10/2017
No. of Questions: (3)
Time: One hour
Using Calculator (No)

University of Palestine

1st. Midterm First Aid
Exam
2017/2018
Total Grade:30 Marks

Instructor Name:
Student No.: _____
Student Name: _____
College Name: تخصص التمريض
Dep. / Specialist: _____
Using Dictionary (No)

Q3- Answer the following questions (15 Marks):

1. Mention the chine of survival?.....(4 mark)
2. Write the steps of CPR for adult victim?..... (8 mark)
3. List the scale of unconsciousness?(4 mark)

**Good luck for all
Dr. Yousif M. Awad**